

# Wet tropospheric corrections comparison : ERA Interim versus ECMWF

Study variable	<b>TRO_HUM_ERA</b>
Reference variable	<b>TRO_HUM_ECMWF</b>
Missions	Topex-Posedon ( <i>tp</i> ), Jason-1 ( <i>j1</i> ), Envisat ( <i>en</i> )
Period	[15705, 21914]

Creation date : 2011/06/16

## Contents

<b>A001</b>	<b>3</b>
<b>A002</b>	<b>6</b>
<b>A003</b>	<b>9</b>
<b>A101</b>	<b>15</b>
<b>A102</b>	<b>18</b>
<b>A103</b>	<b>21</b>
<b>A104</b>	<b>24</b>
<b>A201</b>	<b>27</b>
<b>A202</b>	<b>36</b>
<b>A203</b>	<b>42</b>
<b>A204</b>	<b>51</b>
<b>A206</b>	<b>57</b>

## Study overview

In this study, the wet troposphere model ERA Interim has been compared to the ECMWF wet troposphere correction model.

The impact of using these both wet troposphere corrections on the SSH calculation has been analyzed for Topex-Poseidon, Jason-1 and Envisat missions :

- for Topex-Poseidon : from October 1992 (cycle 4) to October 2006 (Cycle 481)
- for Jason-1 : from January 2002 (cycle 1) to December 2009 (Cycle 294)
- for Envisat : from September 2002 (cycle 9) to December 2009 (Cycle 85)

The ERA Interim wet troposphere correction is based on the ERA INTERIM model which corresponds to the latest global atmospheric reanalysis produced by the European Centre for Medium-Range Weather Forecasts (ECMWF). For more details, see The ERA-Interim reanalysis: configuration and performance of the data assimilation system (Q. J. R. Meteorol. Soc. 137: 553-597, April 2011 A). The reference wet troposphere correction is produced by EUMETSAT which uses 3D data from the ECMWF model to generate wet troposphere values.

All the validation diagnostics displayed in this report has been performed in agreement with the Sea-Level CCI Product Validation Plan (PVP).



## Diagnostic A001 (mission en)

**Name :** Temporal evolution of differences between both altimetric components

**Input data :** Along-track altimetric components

**Description :** The temporal evolution of global statistics (mean, variance, slope) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) . These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses




## Diagnostic A001 (mission j1)

**Name :** Temporal evolution of differences between both altimetric components

**Input data :** Along-track altimetric components

**Description :** The temporal evolution of global statistics (mean, variance, slope) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) . These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses




## Diagnostic A001 (mission tp)

**Name :** Temporal evolution of differences between both altimetric components

**Input data :** Along-track altimetric components

**Description :** The temporal evolution of global statistics (mean, variance, slope) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) . These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses





### Diagnostic A002 (mission en)

**Name :** Map of differences between both altimetric components over all the period

**Input data :** Along-track altimetric components

**Description :** The map of global statistics (mean, standard deviation) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated over a given period which is the longer as possible to have obtain reliable statically results. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses




## Diagnostic A002 (mission j1)

**Name :** Map of differences between both altimetric components over all the period

**Input data :** Along-track altimetric components

**Description :** The map of global statistics (mean, standard deviation) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated over a given period which is the longer as possible to have obtain reliable statically results. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses




## Diagnostic A002 (mission tp)

**Name :** Map of differences between both altimetric components over all the period

**Input data :** Along-track altimetric components

**Description :** The map of global statistics (mean, standard deviation) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated over a given period which is the longer as possible to have obtain reliable statically results. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses




### Diagnostic A003\_a (mission en)

**Name :** Periodogram derived from temporal evolution of altimetric component differences

**Input data :** Along-track altimetric components

**Description :** The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.





## Diagnostic A003\_b (mission en)

**Name :** Periodogram derived from temporal evolution of altimetric component differences

**Input data :** Along-track altimetric components

**Description :** The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Global internal analyses




## Diagnostic A003\_a (mission j1)

**Name :** Periodogram derived from temporal evolution of altimetric component differences

**Input data :** Along-track altimetric components

**Description :** The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Global internal analyses





## Diagnostic A003\_b (mission j1)

**Name :** Periodogram derived from temporal evolution of altimetric component differences

**Input data :** Along-track altimetric components

**Description :** The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Global internal analyses




## Diagnostic A003\_a (mission tp)

**Name :** Periodogram derived from temporal evolution of altimetric component differences

**Input data :** Along-track altimetric components

**Description :** The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Global internal analyses





## Diagnostic A003\_b (mission tp)

**Name :** Periodogram derived from temporal evolution of altimetric component differences

**Input data :** Along-track altimetric components

**Description :** The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Global internal analyses





## Diagnostic A101 (mission en)

**Name :** Temporal evolution of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The temporal evolution of global statistics (mean, standard deviation) of SSH differences are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses




## Diagnostic A101 (mission j1)

**Name :** Temporal evolution of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The temporal evolution of global statistics (mean, standard deviation) of SSH differences are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses




## Diagnostic A101 (mission tp)

**Name :** Temporal evolution of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The temporal evolution of global statistics (mean, standard deviation) of SSH differences are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses




### Diagnostic A102 (mission en)

**Name :** Differences between temporal evolution of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The difference of temporal evolution between the global statistics (mean, standard deviation) of SSH differences are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses




## Diagnostic A102 (mission j1)

**Name :** Differences between temporal evolution of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The difference of temporal evolution between the global statistics (mean, standard deviation) of SSH differences are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses




## Diagnostic A102 (mission tp)

**Name :** Differences between temporal evolution of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The difference of temporal evolution between the global statistics (mean, standard deviation) of SSH differences are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses



### Diagnostic A103 (mission en)


**Name :** Map of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers


**Description :** The differences between maps of SSH crossovers differences (mean, variance) are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses

Mean of SSH with Var\_Stu  
Mission en, cycles 10 to 84



Mean ( cm )  
Mean of SSH with Var\_Ref  
Mission en, cycles 10 to 84



## Diagnostic A103 (mission j1)


**Name :** Map of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers


**Description :** The differences between maps of SSH crossovers differences (mean, variance) are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses

Mean of SSH with Var\_Stu  
Mission j1, cycles 2 to 293



Mean ( cm )  
Mean of SSH with Var\_Ref  
Mission j1, cycles 2 to 293




## Diagnostic A103 (mission tp)

**Name :** Map of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The differences between maps of SSH crossovers differences (mean, variance) are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses



## Diagnostic A104 (mission en)


**Name :** Differences between maps of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The differences between maps of SSH crossovers (derived from diagnostic A103) are calculated from the SSH crossover differences (mean, standard deviation) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses

$\text{VAR}(\text{SSH with Var\_Stu}) - \text{VAR}(\text{SSH with Var\_Ref})$   
Mission en, cycles 10 to 84



## Diagnostic A104 (mission j1)


**Name :** Differences between maps of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The differences between maps of SSH crossovers (derived from diagnostic A103) are calculated from the SSH crossover differences (mean, standard deviation) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses

$\text{VAR}(\text{SSH with Var\_Stu}) - \text{VAR}(\text{SSH with Var\_Ref})$   
Mission j1, cycles 2 to 293



## Diagnostic A104 (mission tp)


**Name :** Differences between maps of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The differences between maps of SSH crossovers (derived from diagnostic A103) are calculated from the SSH crossover differences (mean, standard deviation) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses

$\text{VAR}(\text{SSH with Var\_Stu}) - \text{VAR}(\text{SSH with Var\_Ref})$   
Mission tp, cycles 344 to 480




## Diagnostic A201 a (mission en)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses





## Diagnostic A201\_b (mission en)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses





## Diagnostic A201\_c (mission en)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses




## Diagnostic A201\_a (mission j1)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses





## Diagnostic A201\_b (mission j1)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses




## Diagnostic A201\_c (mission j1)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses




## Diagnostic A201\_a (mission tp)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses





## Diagnostic A201\_b (mission tp)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses




## Diagnostic A201\_c (mission tp)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses




**Diagnostic A202\_a (mission en)**

**Name :** Differences of temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) or separating North and South hemispheres.

Diagnostic type : Global internal analyses





## Diagnostic A202\_b (mission en)

**Name :** Differences of temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) or separating North and South hemispheres.

Diagnostic type : Global internal analyses




## Diagnostic A202\_a (mission j1)

**Name :** Differences of temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) or separating North and South hemispheres.

Diagnostic type : Global internal analyses




## Diagnostic A202\_b (mission j1)

**Name :** Differences of temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) or separating North and South hemispheres.

Diagnostic type : Global internal analyses




## Diagnostic A202\_a (mission tp)

**Name :** Differences of temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) or separating North and South hemispheres.

Diagnostic type : Global internal analyses




## Diagnostic A202\_b (mission tp)

**Name :** Differences of temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) or separating North and South hemispheres.

Diagnostic type : Global internal analyses



### Diagnostic A203\_a (mission en)


**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA


**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses

SLA with Var\_Stu : trends  
Mission en, cycles 10 to 84



Trends (mm/yr)  
SLA with Var\_Ref : trends  
Mission en, cycles 10 to 84



## Diagnostic A203\_b (mission en)


**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA


**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses

SLA with Var\_Stu : trends, even pass numbers  
Mission en, cycles 10 to 84



SLA with Var\_Ref : trends, even pass numbers  
Mission en, cycles 10 to 84



## Diagnostic A203\_c (mission en)


**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA


**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses

SLA with Var\_Stu : trends, odd pass numbers  
Mission en, cycles 10 to 84



SLA with Var\_Ref : trends, odd pass numbers  
Mission en, cycles 10 to 84



## Diagnostic A203\_a (mission j1)


**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA


**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses

SLA with Var\_Stu : trends  
Mission j1, cycles 2 to 293



Trends (mm/yr)  
SLA with Var\_Ref : trends  
Mission j1, cycles 2 to 293





## Diagnostic A203\_b (mission j1)

**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA

**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses





## Diagnostic A203\_c (mission j1)

**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA

**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses





## Diagnostic A203\_a (mission tp)

**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA

**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses





## Diagnostic A203\_b (mission tp)

**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA

**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses



## Diagnostic A203\_c (mission tp)


**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA


**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses

SLA with Var\_Stu : trends, odd pass numbers  
Mission tp, cycles 344 to 480



Trends (mm/yr)  
SLA with Var\_Ref : trends, odd pass numbers  
Mission tp, cycles 344 to 480



**Diagnostic A204 a (mission en)**


**Name :** Differences between maps of SLA

**Input data :** Along track SLA

**Description :** The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Global internal analyses

SLA with Var\_Stu – SLA with Var\_Ref : trends  
Mission en, cycles 10 to 84



## Diagnostic A204\_b (mission en)


**Name :** Differences between maps of SLA

**Input data :** Along track SLA


**Description :** The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Global internal analyses

SLA with Var\_Stu - SLA with Var\_Ref : trends, even pass numbers  
Mission en, cycles 10 to 84



Trends (mm/yr)  
SLA with Var\_Stu - SLA with Var\_Ref : trends, odd pass numbers  
Mission en, cycles 10 to 84



## Diagnostic A204\_a (mission j1)


**Name :** Differences between maps of SLA

**Input data :** Along track SLA

**Description :** The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Global internal analyses

SLA with Var\_Stu – SLA with Var\_Ref : trends  
Mission j1, cycles 2 to 293



## Diagnostic A204\_b (mission j1)


**Name :** Differences between maps of SLA

**Input data :** Along track SLA


**Description :** The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Global internal analyses

SLA with Var\_Stu - SLA with Var\_Ref : trends, even pass numbers  
Mission j1, cycles 2 to 293



Trends (mm/yr)  
SLA with Var\_Stu - SLA with Var\_Ref : trends, odd pass numbers  
Mission j1, cycles 2 to 293



## Diagnostic A204\_a (mission tp)


**Name :** Differences between maps of SLA

**Input data :** Along track SLA

**Description :** The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Global internal analyses

SLA with Var\_Stu – SLA with Var\_Ref : trends  
Mission tp, cycles 344 to 480



## Diagnostic A204\_b (mission tp)


**Name :** Differences between maps of SLA

**Input data :** Along track SLA


**Description :** The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Global internal analyses

SLA with Var\_Stu - SLA with Var\_Ref : trends, even pass numbers  
Mission tp, cycles 344 to 480



Trends (mm/yr)  
SLA with Var\_Stu - SLA with Var\_Ref : trends, odd pass numbers  
Mission tp, cycles 344 to 480




### Diagnostic A206\_a (mission en)

**Name :** Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Global internal analyses



## Diagnostic A206\_b (mission en)


**Name :** Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.


Periodogram of SLA differences, north and south hemispheres (reference period = 1 year)

Mission en, cycles 10 to 84



Periodogram of SLA differences, north and south hemispheres (period = [0, 1 year])

Mission en, cycles 10 to 84





## Diagnostic A206\_a (mission j1)

**Name :** Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Global internal analyses



## Diagnostic A206\_b (mission j1)

**Name :** Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)


**Input data :** Along track SLA

**Description :** The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Global internal analyses


Periodogram of SLA differences, north and south hemispheres (reference period = 1 year)

Mission j1, cycles 2 to 293



Periodogram of SLA differences, north and south hemispheres (period = [0, 1 year])

Mission j1, cycles 2 to 293





## Diagnostic A206\_a (mission tp)

**Name :** Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Global internal analyses



## Diagnostic A206\_b (mission tp)

**Name :** Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)


**Input data :** Along track SLA

**Description :** The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Global internal analyses

Periodogram of SLA differences, north and south hemispheres (reference period = 1 year)

Mission tp, cycles 344 to 480



Periodogram of SLA differences, north and south hemispheres (period = [0, 1 year])

Mission tp, cycles 344 to 480

